

WESTERN ANACAPIA — A SUMMARY OF THE CENOZOIC HISTORY OF THE NORTHERN CHANNEL ISLANDS

D. W. Weaver and D. P. Doerner

University of California, Santa Barbara

Anacapia is a geologic province which forms the southern boundary of the Santa Barbara Embayment and the northern limits of an old, often emergent, land mass - Catalinia. It embraces San Miguel, Santa Rosa, Santa Cruz, and Anacapa islands as well as the western part of the Santa Monica Mountains (Reed and Hollister, 1936).

A study of the geology and evolution of the Northern Channel Islands, lying as they do on the southern edge of the Transverse Ranges (fig. 1) of California, has long held the interest of natural historians, both geologic and biologic. The Transverse Ranges as a whole form an anomalous feature to the general north-south trending structures of western North America. The Murray Escarpment, trending westward 1,900 miles out to sea, is believed to be a structural extension of this continental anomaly. This east-west trend terminates to the east against the great San Andreas fault system and the San Bernardino Range. Not only is it of interest for its obvious structural significance as it is related to one of the great features of the earth's crust, but this transverse feature also has strong biogeographical implications, having roots perhaps as far back as early Mesozoic time. With the progressive provincialization of marine faunas during the Cenozoic (Smith, 1919), these east-west trending mountain ranges, at times partly submerged, partly emerged, served as the limits on more than one occasion for new biogeographic provinces (Kleinpell and Weaver, 1963), some of which became distinctive and conspicuous life areas in the middle and late Cenozoic. Thus the Transverse Ranges serve as a focal point of considerable significance not only for structural but for biogeographic studies as well. In fact, these two fields, tectonics and chorology, are rather intricately interrelated, though perhaps more directly through intermediate climatological phenomena.




FIG. 1.— Diagrammatic relief map to show some important topographic and structural features of Southern California. Modified from Reed & Hollister, 1936

The purpose of this paper is to summarize for the Symposium the paleogeographic implications of the available¹ geologic data for the southern fringe of the Transverse Ranges - the Northern Channel Islands. Realizing that members of the Symposium are primarily interested in Cenozoic migratory land routes to the islands, it should be stated at the outset that present-day Anacapia offers a paucity of evidence for Cenozoic connections to the mainland. In general, ancient migratory routes are best delineated by the land animals and plants of the time. With but two exceptions, the late Pleistocene to sub-Recent dwarf mammoths and the Recent biota of the Northern Channel Islands, such organisms are not known to be preserved. The Cenozoic land masses that are needed for faunal migrations to the islands are inferred from: (1) the presence of continental deposits; (2) the absence of deposits representing major intervals of time, thus permitting the inference that the land was emergent during at least part of the time represented by the hiatus; (3) the fossil and mineralogical composition, texture, and structure of the marine and non-marine sedimentary rocks, which permit ecologic inferences; and (4) the gross structural framework of the province along which any orogenic (mountain building) forces must have operated. The foregoing types of data are available from the literature cited and from recent unpublished work by the authors.

Anacapia is an east-west trending physiographic unit with much of its lower elevations flooded by the Recent seas. Its pre-Tertiary, perhaps Mesozoic, origin is faintly reflected in the metamorphic and granitoid rocks of unknown, but pre-Cenozoic, age which are exposed on Santa Cruz Island and in the Santa Monica Mountains. It is dominated by sedimentary and volcanic rocks of Cretaceous to Miocene age and capped by late Pleistocene terrace formations. Rock exposures on the islands are notable for their complete lack of Pliocene deposits. The pre-Pliocene shales, sandstones, conglomerates, and volcanics are conspicuous for their lateral variation and discontinuity. Areas of post-Eocene erosion and deposition, especially in Catalina, appear to have been uplifted and depressed mainly by block faulting of the basin and range type, i.e., similar to the basins and ranges common in Nevada (fig. 2). Such deformation began in Oligocene time and continued intermittently through the rest of the Cenozoic Era. Many of the blocks reversed their relative vertical positions during this time (Corey, 1954; Emery, 1954).

1. More information is known than is available to the author, since much data involving the off-shore area are of an industrial and thus confidential nature.

as a Miocene land mass with its associated fauna and flora was essentially destroyed in the chalky and siliceous transgressive sea.

Late Miocene time saw a re-emergence of western Anacapia, though the Santa Rosa basin was still evident as a marine trough. However, by Pliocene time even this long persistent topographic low had filled or was uplifted; and emergence was again, as in Oligocene time, general and at a maximum throughout Anacapia. Major basins to the south, undergoing embryonic development between the fault scarps (fig. 2) during Miocene time or earlier, persisted throughout the Pliocene and into the Recent (viz., the Santa Cruz, Santa Monica, and other basins; see fig. 2). The extent of the Pliocene land mass of western Anacapia with its undoubted connections to the mainland must surely have produced a rich land biota. However, no Pliocene marine or non-marine strata are known from any of the Northern Channel Islands; and therefore, no record of the Pliocene life in western Anacapia is preserved. Any deposits that did accumulate were eroded away in the subsequent land environment that persisted along the east-west trending backbone of Anacapia.

Pleistocene and Recent times saw great fluctuations in the sea level; and these, coupled with movements along the established fault systems (fig. 2), all but drowned western Anacapia. Only in late Pleistocene and sub-Recent times did mainland connections sufficiently, although briefly, re-establish themselves to permit a Pleistocene mammalian fauna to reach as far west as San Miguel Island.²

2. In addition to the well known mammoths on Santa Rosa Island, recent work has redocumented the presence of these Pleistocene and sub-Recent dwarfs on San Miguel and Santa Cruz islands. It is also worthy to note that not only did these proboscidians not reach western Anacapia before late Pleistocene because of the lack of an adequate peninsular migration route, but that the elephants as a group did not arrive in the new world until mid-Pleistocene time; and therefore, could not have existed in Anacapia before that time and probably not until post-late Pleistocene terracing, i.e., the sub-Recent or Holocene. The question arises whether sufficient time was then available for dwarfing of the stock. In this connection, it is of considerable interest to note the rate of dwarfism in other mammalian groups as, for example, the dwarfing of Recent horses isolated in the Grand Canyon since their reintroduction by Europeans following the late Pleistocene extinction of *Equus* in North America.

In summary, it can be said that connections during the Cenozoic Era between Anacapia and the mainland were general for the first time in Oligocene time and reached a maximum during the Pliocene and late Pleistocene or sub-Recent, but that the general emergences were in each case separated by an interval of submergence sufficient to all but destroy the biota of the time. Indeed, the middle Miocene transgression is believed by most workers to have completely drowned western Anacapia. Recent studies on Santa Cruz Island, however, suggest that localized remnants of this early Miocene center of volcanic activity may have remained as a vestige of the earlier peninsula. A comparable scene must have existed during Pleistocene time, for marine terraces cover all but the top of Green Mountain on San Miguel and occurred possibly as high as 1,800 feet on the north side of Santa Cruz.

REFERENCES

- Bailey, T. L., and R. H. Jahns. 1954. Geology of the Transverse Range Province, southern California. Calif. Dept. Nat. Resources, Div. Mines Bull., 170:83-106.
- Bremner, C. St. J. 1932. Geology of Santa Cruz Island, Santa Barbara County, California. Santa Barbara Mus. Nat. Hist. Occas. Papers, (1):1-33, 5 pl.
- Bremner, C. St. J. 1933. Geology of San Miguel Island, Santa Barbara County, California. Santa Barbara Mus. Nat. Hist. Occas. Papers, (2):1-23, 4 pl.
- Clements, T. 1955. The Pleistocene history of the Channel Island Region, southern California. *In* Essays in the natural sciences in honor of Captain Allan Hancock, pp. 311-323. Los Angeles.
- Corey, W. H. 1954. Tertiary basins of southern California. Calif. Dept. Nat. Resources, Div. Mines Bull., 170:73-83.
- Emery, K. O. 1954. General geology of the offshore area, southern California. Calif. Dept. Nat. Resources, Div. Mines Bull., 170:107-111.
- Kleinpell, R. M., and D. W. Weaver. 1963. Oligocene biostratigraphy of the Santa Barbara Embayment, California. Univ. Calif. Pub. Geol. Sci., 43:1-250, 38 pl., 8 fig.
- Orr, P. C. 1956. Radiocarbon dates from Santa Rosa Island. I. Santa Barbara Mus. Nat. Hist. Bull., (2):1-10.
- Orr, P. C. 1960. Radiocarbon dates from Santa Rosa Island. II. Santa Barbara Mus. Nat. Hist. Bull., (3):1-10.
- Rand, W. W. 1933. The Geology of Santa Cruz Island, California. Ph.D. thesis. Univ. Calif., Berkeley.

- Reed, R. D., and J. S. Hollister. 1936. Structural evolution of southern California. Tulsa. 157 pp.
- Smith, J. P. 1919. Climatic relations of the Tertiary and Quaternary faunas of the California region. Proc. Calif. Acad. Sci., ser. 4, 9 (4):123-173.
- Smith, W. S. T. 1900. A topographic study of the islands of southern California. Univ. Calif. Pub. Geol. Sci., 2 (7):179-230, 1 pl.
- Stock, C. 1935. Exiled elephants of the Channel Islands, California. Sci. Monthly, 41(3):205-214.
- Vedder, J. G., and R. M. Norris. 1963. Geology of San Nicolas Island, California. U.S. Geol. Surv. Prof. Paper, 369:1-65.